



FRUIT CHART IN RIPENING ORDER

| FRUIT | VARIETY | HEIGHT | POLLINATE |
|--------------------|----------------------------------|--------|------------------|
| Mid June | Blueberry, Northland | 4' | Self |
| Mid June | Gooseberry, Pixwell | 3-4' | Self |
| June | Cherry, 4-N-1 Combo (Semi-Dwarf) | 12-15' | Self |
| June | Cherry, Sweet | 12-15' | Needs Pollinator |
| June | Cherry, Tart pie | 12-15' | Self |
| Mid Summer | Blackberry, Black Satin | 5-6' | Self |
| Mid Summer | Boysenberry, Thornless | 5-6' | Self |
| Mid Summer | Raspberry, Canby Red | 5-6' | Self |
| July-August | Peach | 12-15' | Self |
| Mid August | Plum, Santa Rosa (Standard) | 15-18' | Self |
| August-September | Grape, Concord | 20-25' | Self |
| August-First Frost | Pear, 4-N-1 Combo (Semi-Dwarf) | 12-15' | Self |
| August-First Frost | Pear, Single variety | 12-15' | Needs Pollinator |
| Early September | Plum, Stanley Prune-Plum (Dwarf) | 8-10' | Self |
| Mid September | Grape, Niagara | 20-25' | Self |
| September | Apple 4-N-1 Combo (Dwarf) | 8-10' | Self |
| September | Apple Single variety | 12-15' | Needs Pollinator |

FRUIT TREE SPRAY SCHEDULE

| | |
|---|--|
| 1. PRE-PINK STAGE (just before buds swell) | ferti•lome Horticultural Oil Spray Kills overwintering stages of insects and mites. ferti•lome Broad Spectrum Fungicide Prevents Peach Leaf Curl. |
| 2. PETAL FALL (most petals have fallen) | ferti•lome Fruit Tree Spray Spray every week until fruit shows. ferti•lome Spinosad Worm killer, treat every 2 weeks until harvest. |
| 3. COVER SPRAYS (after fruit set) | ferti•lome Fruit Tree Spray Every week for four weeks, then every two weeks until harvest. |
| 4. DORMANT SPRAY (after leaves fall) | ferti•lome Horticultural Oil Spray Kills overwintering stages of insects and mites. |

Revised 3/2025